EXTRA STRENGTH PAIN RELIEVER- acetaminophen, aspirin and caffeine tablet, coated Chain Drug Consortium, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

premier value 226

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain Reliever

Pain reliever aid

Use

• temporarily relieves minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use

this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these

symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- · more than directed
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher

if you

are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartheat.

Do not use

- if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug

contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking

- any other drug containing an NSAID (prescription or nonprescription)
- a blood thinning (anticoagulant) or steroid drug
- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts more than 3 days
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not

to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose Warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12: ask a doctor

Other Information

- store at controlled room temperature 15° 25°C (59°-77°F)
- read all product information before using

Inactive ingredients

cellulose, fractionated coconut oil, hypromellose, magnesium silicate, maltodextrin, polydextrose, povidone, silica, starch, stearic acid, titanium dioxide.

Package label

NDC 68016-110-00

*Compare to the active ingredients in Excedrin

Premier Value

Extra Strength

PAIN RELIEVER

Acetaminophen 250mg

Aspirin 250mg

Caffeine 65mg

100 Caplets



Drug Facts (continued)

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervounses, irritability, steeplessness, and, occasionally, rapid heartbeat,

Do not use I if you have ever had an allergic reaction to pirin or any other pain reliever/fever reducer

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether drug contains acetaminophen, ask a doctor or pharmacist,

- and command advantagement and a doctor or pharmacie.

 Ask a doctor before use if
 you have liver disease
 stomach bleeding warning applies to you
 you have a history of stomach problems, such as heartbun
 you have high blood pressure, heart disease, liver cirrhosis
- you are taking a diuretic you have asthma.
- Ask a doctor or pharmacist before use if you are taking: any other drug containing an NSAID (prescription or
- nonprescription)

 a blood thinning (anticoagulant) or steroid drug

 a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serior condition

- Stop use and ask a doctor if

 a nallergic reaction occurs, Seek medical help right sway,
 you experience any of the following signs of stomach
 bleeding

 fiel faint = womit blood

 have bloody or black stocks

- In feel tent III womit topod
 have bloody or black stocks
 have stomach pain that does not get better
 inging in the ears or less of hearing occurs
 painful area is red or swelling
 pain gets worse or less for more than 10 days
 fever gets worst or less for more than 3 days
 area one symptoms appear

- any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional ore use. It is especially important not to use aspirin during before use. It is especially important not to use asymm during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children.

Overdose Warning: Taking more than the recommended does (overdose) may cause their damage, in case of overdos get medical help or contact a Poison Control Center right aware. Duties medical statements.

away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- drinks full glass of water with each dose
 adults and children 12 years and over: take 2 caplets
 children under 12: ask a doctor

Other information

■ store at controlled room temperature 15* - 25*C

(59° - 77°F) read all product information before using

Inactive ingredients cellulose, fractionated cochnut of, hypromellose, magnesium silicate, majlodextrin, polydextrose, povidone, silica, starch, stearic acid, titanium

*This product is not manufactured or distributed by the owner of the registered trademark Excedrin®.



SAVE CARTON FOR COMPLETE DRUG FACTS

Drug Facts

Purposes

Active ingredients (in each caplet)

Pain reliever air

nonsteroidal anti-inflammatory drug Use

■ temporarily relieves minor aches and pains

Caffeine 65 mg.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product, When using this product, it changes in behavior with nausea and voreiting occur, consult a dictor because these symptoms could be an early sign of Reye's syndrome, a rare but serious lithess.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- which may include:

 hives If facial swelling

 asthma (wheezing) If shock
 Liver warning: This product contains scetaminophen.
- Severe liver damage may occur if you take

 more than directed

 with other drugs containing acetaminophen

 3 or more alcoholic drinks every day while using this

product
Stomach bleeding warning: This product contains an
NSAID, which may cause severe stomach bleeding. The
chance is higher if you
are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
 take other drugs containing prescription or nonprescript
 NSAIDs (aspirin, ibuproten, naproxen, or others)
 have 3 or more alcoholic drinks every day while using
- this product
- more or for a longer time than directed



REV GC226-1111

ORAL

EXTRA STRENGTH PAIN RELIEVER

acetaminophen, aspirin, caffeine tablet, coated

Product Information

Route of Administration

Product Type HUMAN OTC DRUG Item Code (Source) NDC:68016-110

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6 A5W338 E) (CAFFEINE - UNII:3G6 A5W338 E)	CAFFEINE	65 mg

DEA Schedule

Inactive Ingredients	
Ingredient Name	Strength

MAGNESIUM SILICATE (UNII: 9B9691B2N9)	
MALTO DEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
PO VIDO NES (UNII: FZ989GH94E)	
SILICON DIO XIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
HYPROMELLOSES (UNII: 3NXW29 V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics			
Color	white (WHITE)	Score	no score
Shape	CAPSULE (Capsule)	Size	17mm
Flavor		Imprint Code	GC226
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:68016-110-00	1 in 1 CARTON			
1		100 in 1 BOTTLE			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/01/2008	

Labeler - Chain Drug Consortium, LLC (101668460)

Registrant - Geri-Care Pharmaceutical Corp (611196254)

Establishment			
Name	Address	ID/FEI	Business Operations
Geri-Care Pharmaceutical Corp		611196254	repack(68016-110)

Revised: 11/2014 Chain Drug Consortium, LLC